


STRANGULATION

Visible Signs (may not be present)



Additional Signs and Symptoms

A larger version of the graphic above which contains detailed signs and symptoms is available for download at strangulationtraininginstitute.com/resources/library/pediatric/

This project is supported all or in part by Grant No. 2016-TA-AX-K067 awarded by the Office on Violence Against Women, U.S. Department of Justice. The opinions, findings, conclusions, and recommendations expressed in this publication/program/exhibition are those of the author(s) and do not necessarily reflect the views of the Department of Justice, Office on Violence Against Women.

This printing of this brochure is supported by subgrant number 20-VOCA-46 awarded through the Federal Office for Victims of Crime as administered by the Kansas Governor's Grants Program. The opinions, findings, conclusions, or recommendations expressed in this publication, program, or exhibition are those of the author(s) and do not necessarily reflect the views of the Office of the Kansas Governor or the U.S. Department of Justice. This program does not discriminate on the basis of race, color, religion, national origin, sex, age, or disability.

Strangulation is often under-recognized in children but no less serious than in adults. Unconsciousness may happen within seconds and death within minutes. Children may be strangled when caregivers lose control, as part of physical and/or sexual assault, or as a way of demonstrating ultimate power and control over the child. Regardless, strangulation of a child can have long-lasting physical and mental health effects and can result in death even months later.

Child victims of strangulation may feel terror and extreme pain. If strangulation continues, unconsciousness will follow. Before sliding into unconsciousness, a child victim may resist violently, producing injuries to their own neck or to the face or hands of their attacker. These defensive injuries may not be present in young or developmentally disabled children, or if the victim is physically or chemically restrained.

Observing Changes

Documentation by photographs organized in order, for a period of days after the attack is very helpful in beginning and building a journal of proof. Victims should be given medical attention if they experience difficulty breathing, speaking, swallowing or experience nausea, vomiting, lightheadedness, headache or holding head, accidental urination and/or bowel movement in children not diapered. A medical evaluation may be extremely important in detecting internal injuries and saving a life.

Losing Consciousness

Victims may lose awareness or faint by any one or all of the following methods: blocking of the blood vessels from the heart in the neck (taking away oxygen from the brain), blocking of the large veins in the neck (preventing deoxygenated blood from exiting the brain), and closing off the tube from the mouth to the lungs, making breathing impossible.



Notify police, 911, and medical personnel immediately

Wichita Police Department

455 N. Main
Wichita, KS 67202

316-268-4156

www.wichitapolice.com



ALLIANCE for
HOPE
INTERNATIONAL
a program of

strangulationtraininginstitute.com

Graphic Design and Illustrations by Yesenia Aceves

STRANGULATION
What Parents and Caregivers
NEED TO KNOW

Monitor the Child's SIGNS

Date & Time Photograph and Journal the Child's Signs

| | |
|--|--|
| | |
| | |
| | |
| | |
| | |
| | |

Monitor the Child's SYMPTOMS

Date & Time Journal the Child's Symptoms

| | |
|--|--|
| | |
| | |
| | |
| | |
| | |
| | |

Date & Time Journal Any Other Sensation

| | |
|--|--|
| | |
| | |

Signs of Strangulation

Head- pinpoint red spots (petechiae) on scalp, hair pulled, bump(s), skull fracture(crack), concussion.

Face- red or flushed, petechiae, scratch marks.

Eyes and Eyelids- petechiae to the left or right eyeball, bloodshot eyes.

Ear- petechiae (external and/or ear canal), bleeding from ear canal.

Nose- bloody nose, broken nose, petechiae.

Mouth- bruising, swollen tongue, swollen lips, cuts/abrasions(scrapes).

Under the chin- redness, scratch marks, bruise(s), abrasions.

Neck- redness, scratch marks, fingernail marks, bruise(s), abrasions, swelling, ligature(tie) or clothing marks.

Chest and Shoulders- redness, scratch marks, bruise(s), abrasions.

Symptoms of Strangulation

Voice changes- raspy and/or hoarse voice, coughing, unable to speak, complete loss of voice.

Swallowing changes- trouble swallowing, painful swallowing, neck pain, nausea/vomiting, drooling.

Breathing changes- difficulty breathing, hyperventilation, unable to breathe.

Behavioral changes- restlessness or aggressiveness, problems concentrating, amnesia(loss of memory), agitation, Post-traumatic Stress Syndrome, hallucinations.

Vision changes- complete loss or black & white vision, seeing 'stars', blurry, darkness, fuzzy around the eyes.


Hearing changes- complete loss of hearing, gurgling, ringing, buzzing, popping, pressure, tunnel-like hearing.

Other changes- Memory loss, unconsciousness, dizziness, headaches, involuntary urination or bowel movement in potty-trained child, loss of strength, going limp.


Diagrams to Mark Visible Injuries

Use a pen or a marker to indicate any visible signs and/or symptoms.


Front




Under Chin




Right Side



Left Side



Back



Inside Mouth

